	Weight Bearing	<u>Brace</u>	ROM	<u>Exercises</u>
Phase I 0-2 weeks	Heel touch only.	0-2 weeks: Locked in full extension at all times.  Off for hygiene and exercises only.	Gentle 0-90° as tolerated for first 2 weeks.	<b>0-2 weeks</b> : Quad sets, SLR in brace, calf pumps, passive leg hangs to 90°, modalities.
Phase II 2-6 weeks	Heel touch only.	Unlocked during the day 0-90°, off at night. Off for hygiene, rest, and exercises.	Maintain full extension and progress to full flexion as tolerated.	PROM/AAROM to tolerance, patella and tibiofibular joint mobs.  Begin floor based core and glute/ quad/hamstring exercises.
Phase III 6-8 weeks	Advance 25% weekly until full.		Full.	6-8 weeks: Advance closed chain exercises, gait normalization, eccentric quads, eccentric hamstrings; advance core, glutes, and pelvic stability.  Begin stationary bike as tolerated after 6 weeks.  8-12 weeks: Advance hamstring work, proprioception/balance exercises; hip/core/glutes.
Phase IV 8-16 weeks	Full.	None.	Full.	Progress Phase III exercises and functional activities: walking lunges, planks, bridges, swiss ball, half-bosu exercises.  Advance core/glutes, balance, flexibility.  May advance to elliptical, pool as tolerated after 12 weeks.
Phase V >16 weeks	Full.	None.	Full.	Slowly advance all activity as tolerated.  Progress to functional training, including running, jumping, pivoting, and sports after 16 weeks.

<sup>\*</sup>Brace may be removed at night after first post-operative visit (day 7-14) if directed.