

	<u>Weight Bearing</u>	<u>Brace</u>	<u>ROM</u>	<u>Exercises</u>
<u>Phase I</u> <u>0-8 weeks</u>	Full with brace.	0-2 weeks: Locked in full extension at all times.* Off for CPM, hygiene, and exercises only. 2-4 weeks: Unlocked 0-45°, worn daytime only. 4-6 weeks: Unlocked 0-90°, worn daytime only. Discontinue at 6 weeks.	0-2 weeks: Gentle passive 0-45° as tolerated. 2 weeks+: advance as tolerated. 0-6 weeks: use CPM 6 hours/day, advance 5-10° per day as tolerated, d/c at 6 weeks. <u>Goals:</u> 0-2 wks: 0-30° 2-4 wks: 0-60° 4-6 wks: 0-90°	0-2 weeks: Quad sets, SLR, calf pumps, passive leg hangs to 45° at home. 2-6 weeks: PROM/AAROM to tolerance, patella and tibiofibular joint mobs, quad, hamstring, and glute sets, SLR, side-lying hip and core. 6-8 weeks: Addition of heel raises, closed chain exercises, gait normalization, eccentric quads, eccentric hamstrings; advance core, glutes, and pelvic stability.
<u>Phase II</u> <u>8-12 weeks</u>	Full.	None.	Full.	Progress closed chain activities and gait training. Begin unilateral stance activities, balance training, hamstring work, hip/core/glutes. Begin stationary bike at 10 weeks with low resistance as tolerated.
<u>Phase III</u> <u>12-24 weeks</u>	Full.	None.	Full.	Advance Phase II exercises; maximize core/glutes, pelvic stability work, eccentric hamstrings, balance. Swimming as tolerated after 12 wks. Elliptical as tolerated after 16 wks.
<u>Phase IV</u> <u>6-12 months</u>	Full.	None.	Full.	Advance functional activity without impact. Return to sport-specific activity and impact when cleared by physician after 6-8 months.

*Brace may be removed at night after first post-operative visit (day 7-14) if directed.

*Alignment correction procedures do not change protocol (ex: HTO, DFO, TTO).